

Effective Supergravity from Heterotic M–Theory and its Phenomenological Implications

Carlos Muñoz*

*Departamento de Física Teórica C-XI and Instituto de Física Teórica C-XVI
Universidad Autónoma de Madrid, 28049 Cantoblanco, Madrid, Spain
E-mail: carlos.munnoz@uam.es*

ABSTRACT: In this talk I summarize several recent results concerning the four–dimensional effective supergravity obtained using a Calabi–Yau compactification of the $E_8 \times E_8$ heterotic string from M–theory. A simple macroscopic study is provided expanding the theory in powers of two dimensionless variables. Higher order terms in the Kähler potential are identified and matched with the heterotic string corrections. In the context of this M–theory expansion, I discuss several phenomenological issues: universality of soft scalar masses, relations between the different scales of the theory (eleven–dimensional Planck mass, compactification scale and orbifold scale) in order to obtain unification at 3×10^{16} GeV or lower values, soft supersymmetry–breaking terms, and finally charge and colour breaking minima. The above analyses are also carried out in the presence of (non–perturbative) five–branes.

FTUAM 99/19
IFT-UAM/CSIC-99-24
June 1999

1. Introduction and summary

One of the most exciting proposals of the last years in string theory, consists of the possibility that the five distinct superstring theories in ten dimensions plus supergravity in eleven dimensions be different vacua in the moduli space of a single underlying eleven–dimensional theory, the so–called M–theory [1]. In this respect, Hořava and Witten proposed that the strong–coupling limit of $E_8 \times E_8$ heterotic string theory can be obtained from M–theory. They used the low–energy limit of M–theory, eleven–dimensional supergravity, on a manifold with boundary (a S^1/Z_2 orbifold), with the E_8 gauge multiplets at each of the two ten–dimensional boundaries (the orbifold fixed planes) [2].

In the present paper I will summarize several recent results concerning the four–dimensional

implications of this so called heterotic M–theory. In particular, I will concentrate on the analysis of the effective supergravity obtained by compactifying heterotic M–theory on a six–dimensional Calabi–Yau manifold and its phenomenological consequences.

The effective action of this limit has been systematically analyzed in an expansion in powers of $\kappa^{2/3}$, where κ^2 denotes the eleven–dimensional gravitational coupling [3]. As was noticed in [4], this leads to an expansion parameter which scales as $\kappa^{2/3} \rho / V^{2/3}$, where $\pi \rho$ denotes the length of the eleventh segment and V is the Calabi–Yau volume. At the leading order in this expansion, the Kähler potential, superpotential and gauge kinetic functions have been computed in [4, 5, 6, 7]. It is rather easy to determine the order ϵ_1 correction to the leading order gauge kinetic functions [4, 8, 7, 9], while it is much more nontrivial to compute the order ϵ_1 correction to the leading order Kähler potential, which was recently

*I thank K.Choi, H.B. Kim and D.G. Cerdeño for their collaboration in this project.

done by Lukas, Ovrut and Waldram [10]. It was argued in [11] that the holomorphy and Peccei–Quinn symmetries guarantee that there is *no* further correction to the gauge kinetic functions and the superpotential at any finite order in the M–theory expansion, similarly to the case of the perturbative heterotic string [12].

On the other hand, as is well known, the four–dimensional effective action of the weakly coupled heterotic string theory can be expanded in powers of the two dimensionless variables: the string coupling $\epsilon_s = e^{2\phi}/(2\pi)^5$ and the world-sheet sigma–model coupling $\epsilon_\sigma = 4\pi\alpha'/V^{1/3}$. It was suggested in [11] that the effective action of M–theory can be similarly analyzed by expanding it in powers of the two dimensionless variables: $\epsilon_1 = \kappa^{2/3}\pi\rho/V^{2/3}$ and $\epsilon_2 = \kappa^{2/3}/\pi\rho V^{1/3}$. The latter is the straightforward generalization of the string world–sheet coupling $\sim \alpha'/V^{1/3}$ to the membrane world–volume coupling $\sim \kappa^{2/3}/\rho V^{1/3}$ since $\kappa^{2/3}$ may be identified as the inverse of the membrane tension. Note that in the M–theory limit, heterotic string corresponds to a membrane stretched along the eleventh dimension. In this framework the Kähler potential is expected to receive corrections which are higher order in ϵ_1 or ϵ_2 . An explicit computation of these higher order corrections will be highly non-trivial since first of all the eleven–dimensional action is known only up to the terms of order $\kappa^{2/3}$ relative to the zeroth order action (except for the order $\kappa^{4/3}$ four-gaugino term) and secondly the higher order computation of the compactification solution and its Kaluza-Klein reduction are much more complicated.

In section 2, we will provide a simple macroscopic analysis of the four–dimensional effective supergravity action by expanding it in powers of ϵ_1 and ϵ_2 . Possible higher order corrections in the Kähler potential are identified and matched with the heterotic string corrections, and their size is estimated for the physically interesting values of moduli [11]. The validity of this procedure has been explicitly checked in [15] in the case of M–theory compactified on $S^1/Z_2 \times K3 \times T^2$.

On the other hand, we will also discuss in detail how these effective supergravity models can be strongly constrained by imposing the phenomenological requirement of universal soft scalar

masses, in order to avoid dangerous flavour changing neutral current phenomena. As pointed out in [13], there is a simple solution to avoid this problem: to work with Calabi–Yau spaces with one Kähler modulus T only. Of course, the existence of such spaces, as e.g. the quintic hypersurface in CP^4 , and their universality properties was also known in the context of the weakly–coupled heterotic string, however the novel fact in heterotic M–theory, is that model building is relatively easy. For example, in the presence of non–standard embedding and five–branes (non–perturbative objects located at points throughout the orbifold interval [3]) to obtain three–generation models with realistic gauge groups, as for example $SU(5)$, is not specially difficult [14].

Other phenomenological implications of heterotic M–theory, turn out to be also advantageous with respect to the ones of the perturbative heterotic–string theory. First of all, the resulting four–dimensional effective theory can reconcile the observed Planck scale $M_{Planck} = 1.2 \times 10^{19}$ GeV with the phenomenologically favored GUT scale $M_{GUT} \approx 3 \times 10^{16}$ GeV in a natural manner, providing an attractive framework for the unification of couplings [3, 4]. This is to be compared to the weakly–coupled heterotic string where $M_{string} \approx 8.5 \times 10^{17}$ GeV. Another phenomenological virtue of the M–theory limit is that there can be a QCD axion whose high energy axion potential is suppressed enough so that the strong CP problem can be solved by the axion mechanism [4, 8]. About the issue of supersymmetry breaking, the possibility of generating it by the gaugino condensation on the hidden boundary has been studied [16, 7, 17, 18, 19, 20] and also some interesting features of the resulting soft supersymmetry–breaking terms were discussed. In particular, gaugino masses turn out to be of the same order as squark masses [7]. This is welcome since gaugino masses much smaller than squark masses, as in the weakly–coupled heterotic string case, may give rise to a hierarchy problem [21]. For example, the experimental lower bound on gluino masses of 150 GeV would imply scalar masses larger than 1 TeV. Besides, the phenomenologically favored vacuum expectation values of the moduli can be obtained with several gaugino condensates with the appropri-

ate hidden matter [20], similarly to the case of the weakly-coupled heterotic string [22]. However, it is fair to say that unlike the latter non-perturbative membrane instantons are also necessary in M-theory to obtain the desired minimum.

Several of the above mentioned phenomenological issues will be analyzed in the next sections. In section 3 we will concentrate in the case of standard and non-standard embedding vacua [23, 24, 25, 26], whereas in section 4 vacua in the presence of five-branes [26, 13] are studied. The latter are characterized basically by new moduli Z_n associated with the five-brane positions in the orbifold dimension. In both cases we will perform a detailed study of the different scales of the theory, as well as a systematic analysis of the soft supersymmetry-breaking terms.

Concerning the former, the relations between the eleven-dimensional Planck mass, the Calabi-Yau compactification scale and the orbifold scale, taking into account higher order corrections to the leading order formulae, will be analyzed [27]. Identifying the compactification scale with the GUT scale, to obtain $M_{GUT} \approx 3 \times 10^{16}$ GeV is simpler in non-standard embedding models than in standard ones. In the presence of five-branes, M_{GUT} can be obtained more easily. On the other hand, going away from perturbative vacua, it was recently realized that the string scale may be anywhere between the weak scale and the Planck scale [28] and the size of the extra dimensions may be as large as a millimetre [29]. Whether or not all these scenarios¹ are possible in the context of heterotic M-theory has been analyzed recently in [30] with interesting results: to lower the unification scale (and therefore the eleven-dimensional Planck scale which is around two times bigger) to intermediate values $\approx 10^{11}$ GeV or 1 TeV values or to obtain the radius of the orbifold as large as a millimetre is in principle possible in some special limits. However, it has been pointed out in [27] that the necessity of a fine-tuning or the existence of a hierarchy problem renders these possibilities unnatural. Although

¹To trust them would imply to assume that Nature is trying to mislead us with an apparent gauge coupling unification at M_{GUT} . In this sense, a reasonable doubt about those possibilities is healthy.

new possibilities arise in the presence of five-branes in order to lower the scales, again at the cost of introducing a huge hierarchy problem.

We will also analyze the soft supersymmetry breaking terms under the general assumption that supersymmetry is spontaneously broken by the auxiliary components of the bulk moduli superfields (dilaton S and modulus T) [11, 18, 31, 27]. It is examined in particular how the soft terms vary when one moves from the weakly-coupled heterotic string limit to the strongly-coupled limit. The presence of new parameters in the formulae gives rise to different pattern of soft terms. This is also the case of models with five-branes where at least a new goldstino angle, associated to a modulus Z_1 , must be included in the computation [13, 32, 27]. Unlike the weakly-coupled case, scalar masses larger than gaugino masses can be obtained [27]. Low-energy ($\approx M_W$) sparticle spectra [33, 11, 34, 35, 27] are also discussed.

Finally, the existence of charge and colour breaking minima is analyzed. As is well known, the presence of scalar fields with colour and electric charge in supersymmetric theories induces the possible existence of dangerous charge and colour breaking minima, which would make the standard vacuum unstable [36]. They impose very strong constraints on supergravity models from heterotic M-theory [37, 38]. In particular, standard-embedding models turn out to be excluded on these grounds, similarly to the perturbative heterotic-string situation [38]. Possible solutions to this problem are discussed.

2. Four-dimensional effective supergravity

Here we will analyze the four-dimensional effective supergravity obtained by compactifying heterotic M-theory on a six-dimensional Calabi-Yau manifold.

2.1 Expansions

Let us first discuss possible perturbative expansions of the four-dimensional effective supergravity. As in the case of the weakly coupled heterotic string theory, the effective supergravity of com-

pactified M-theory contains two model-independent moduli superfields S and T whose scalar components can be identified as

$$\begin{aligned} \text{Re}(S) &= \frac{1}{2\pi(4\pi)^{2/3}} M_{11}^6 V, \\ \text{Re}(T) &= \frac{6^{1/3}}{2(4\pi)^{4/3}} M_{11}^3 V^{1/3} \pi \rho, \end{aligned} \quad (2.1)$$

with M_{11} denoting the eleven-dimensional Planck mass, $\kappa^2 = M_{11}^{-9}$. The above normalizations of S and T have been chosen to keep the conventional form of the gauge kinetic functions in the effective supergravity. (See (2.6) for our form of the gauge kinetic functions. Our S and T correspond to $\frac{1}{4\pi}S$ and $\frac{1}{8\pi}T$ of [8] respectively.)

The moduli S and T can be used to define various kind of expansions which may be applied for the low-energy effective action. For instance, in the weakly coupled heterotic string limit, we have

$$\begin{aligned} \text{Re}(S) &= e^{-2\phi} \frac{V}{(2\alpha')^3}, \\ \text{Re}(T) &= \frac{6^{1/3}}{32\pi^3} \frac{V^{1/3}}{2\alpha'}, \end{aligned} \quad (2.2)$$

where ϕ and $\sqrt{2\alpha'}$ denote the heterotic string dilaton and length scale respectively. One may then expand the effective action of the heterotic string theory in powers of the string loop expansion parameter ϵ_s and the world-sheet sigma model expansion parameter ϵ_σ :

$$\begin{aligned} \epsilon_s &= \frac{e^{2\phi}}{(2\pi)^5} \approx 0.3 \frac{[4\pi^2 \text{Re}(T)]^3}{\text{Re}(S)}, \\ \epsilon_\sigma &= \frac{4\pi\alpha'}{V^{1/3}} \approx 0.5 \frac{1}{4\pi^2 \text{Re}(T)}. \end{aligned} \quad (2.3)$$

Here we are interested in the possible expansion in the M-theory limit of the strong heterotic-string coupling $\epsilon_s \gg 1$ for which $\pi\rho \gtrsim M_{11}^{-1}$ and $V \gtrsim M_{11}^{-6}$ and so the physics can be described by eleven-dimensional supergravity. Since we have two independent length scales, ρ and $V^{1/6}$, there can be two dimensionless expansion parameters in the M-theory limit also. As discussed in the introduction there are two natural candidates, ϵ_1 and ϵ_2 , to be the expansion parameters of the four-dimensional effective supergravity action of the Hořava-Witten M-theory. Using $\kappa^2 = M_{11}^{-9}$

these can be written as

$$\begin{aligned} \epsilon_1 &= \frac{\pi\rho}{M_{11}^3 V^{2/3}} \approx \frac{\text{Re}(T)}{\text{Re}(S)}, \\ \epsilon_2 &= \frac{1}{M_{11}^3 \pi\rho V^{1/3}} \approx \frac{1}{4\pi^2 \text{Re}(T)}, \end{aligned} \quad (2.4)$$

where (2.1) has been used to arrive at this expression of ϵ_1 and ϵ_2 . Note that $\epsilon_1 \epsilon_2 \approx 1/[4\pi^2 \text{Re}(S)] \approx \alpha_{GUT}/\pi$ which is essentially the four dimensional field theory expansion parameter. Thus if one goes to the limit in which one expansion works better while keeping the realistic value of α_{GUT} , the other expansion becomes worse. Here we will simply assume that both ϵ_1 and ϵ_2 are small enough so that the double expansion in ϵ_1 and ϵ_2 provides a good perturbative scheme for the effective action of M-theory. As we will see later, it turns out that this expansion works well even when ϵ_1 becomes of order one, which is in fact necessary to have $M_{GUT} \approx 3 \times 10^{16}$ GeV.

To be explicit, let us consider a simple compactification on a Calabi-Yau manifold with the Hodge-Betti number $h_{1,1} = 1$. In this model, the low-energy degrees of freedom include first the gravity multiplet and S and T which are the massless modes of the eleven-dimensional bulk fields. We also have gauge and charged matter superfields associated to the observable and hidden sector gauge groups, $G_O \times G_H \subset E_8 \times E_8$, where $G_O(G_H)$ is located at the boundary $x^{11} = 0(x^{11} = \pi\rho)$ with x^{11} denoting the orbifold coordinate. From now on, we will use as our notation the subscript $O(H)$ for quantities and functions of the observable(hidden) sector.

It is then easy to compute the Kähler potential K , the observable and hidden sector gauge kinetic functions f_O and f_H , and the superpotential W at the leading order in the M-theory expansion. Obviously the leading contribution to the moduli Kähler metric is from the eleven-dimensional bulk field action which is of order κ^{-2} , while the charged matter Kähler metric, the gauge kinetic functions, and the charged matter superpotential receive the leading contributions from the ten-dimensional boundary action which is of order $\kappa^{-4/3}$. One finds [4, 5, 6, 7]

$$\begin{aligned} K &= -\ln(S + \bar{S}) - 3\ln(T + \bar{T}) + \frac{3}{T + \bar{T}} C^p \bar{C}^p \\ f_O &= f_H = S, \end{aligned}$$

$$W = d_{pqr} C^p C^q C^r, \quad (2.5)$$

where d_{pqr} are constant coefficients and C^p are the matter fields, i.e. the effective supergravity computed at the leading order in the M-theory expansion is the same as the effective supergravity of the weakly-coupled heterotic string computed at the leading order in the string loop and sigma model perturbation theory.

The holomorphy and the Peccei-Quinn symmetries imply that there is no correction to the superpotential at any finite order in the S and T -dependent expansion parameters ϵ_1 and ϵ_2 . However the gauge kinetic functions can receive a correction at order ϵ_1 in a way consistent with the holomorphy and the Peccei-Quinn symmetries. This correction can be determined by a direct M-theory computation [3] or by matching the string loop threshold correction to the gauge kinetic function [4, 8, 9, 7]. The result is

$$f_O = S + \beta_O T, \quad f_H = S + \beta_H T, \quad (2.6)$$

where the model-dependent integer coefficients $\beta_{O,H} = \frac{1}{8\pi^2} \int \omega \wedge [\text{tr}(F_{O,H} \wedge F_{O,H}) - \frac{1}{2} \text{tr}(R \wedge R)]$, for the Kähler form ω normalized as the generator of the integer (1,1) cohomology², and they fulfil the following condition:

$$\beta_O + \beta_H = 0, \quad (2.7)$$

with β_O always positive in the case of the standard embedding of the spin connection into one of the E_8 gauge groups. Positive and negative values are possible for non-standard embedding cases [24, 25, 26].

Let us now consider the possible higher order corrections to the Kähler potential. With the Peccei-Quinn symmetries, the Kähler potential can be written as $K = \hat{K}(S + \bar{S}, T + \bar{T}) + Z(S + \bar{S}, T + \bar{T}) C^p \bar{C}^p$ with $\hat{K} = \hat{K}_0 + \delta\hat{K}$, $Z = Z_0 + \delta Z$. Here $\hat{K}_0 = -\ln(S + \bar{S}) - 3\ln(T + \bar{T})$ and $Z_0 = 3/(T + \bar{T})$ denote the leading order results in (2.5), while $\delta\hat{K}$ and δZ are the higher order corrections. Before going to the M-theory expansion of $\delta\hat{K}$ and δZ , it is useful to note that the bulk physics become blind to the existence of

²Usually β is considered to be an arbitrary real number. For T normalized as (2.1), it is required to be an integer [8].

boundaries in the limit $\rho \rightarrow \infty$. However some of the boundary physics, e.g. the boundary Calabi-Yau volume, can be affected by the integral of the bulk variables over the eleventh dimension and then they can include a piece linear in ρ [3]. This implies that $\delta\hat{K}/\hat{K}_0$, being the correction to the pure bulk dynamics, contains only a non-negative power of $1/\rho$ in the M-theory expansion, while $\delta Z/Z_0$ which concerns the couplings between the bulk and boundary fields can include a piece linear in ρ . Since $\epsilon_1^n \epsilon_2^m \sim \rho^{n-m}$, one needs $m \geq n$ for the expansion of $\delta\hat{K}/\hat{K}_0$ and $m \geq n-1$ for the expansion of $\delta Z/Z_0$. Taking account of these, the M-theory expansion of the Kähler potential is given by [11]

$$\begin{aligned} \delta\hat{K} &= \sum_{(n+m \geq 1, m \geq n)} A_{nm} \epsilon_1^n \epsilon_2^m = \sum_{m \geq 1} \frac{A_{0m}}{[4\pi^2 \text{Re}(T)]^m} \\ &\quad + \frac{A_{11}}{4\pi^2 \text{Re}(S)} \left[1 + \mathcal{O}\left(\frac{1}{4\pi^2 \text{Re}(S)}, \frac{1}{4\pi^2 \text{Re}(T)}\right) \right] \\ \delta Z &= \frac{3}{(T + \bar{T})} \sum_{(n+m \geq 1, m \geq n-1)} B_{nm} \epsilon_1^n \epsilon_2^m \\ &= \frac{3}{(T + \bar{T})} \sum_{m \geq 1} \frac{B_{0m}}{[4\pi^2 \text{Re}(T)]^m} + \frac{3B_{10}}{2\text{Re}(S)} \\ &\quad \times \left[1 + \mathcal{O}\left(\frac{1}{4\pi^2 \text{Re}(S)}, \frac{1}{4\pi^2 \text{Re}(T)}\right) \right] \end{aligned} \quad (2.8)$$

where the $n = 0$ terms are separated from the other terms with $n \geq 1$.

The above expansion would work well in the M-theory limit: $[4\pi^2 \text{Re}(T)]^3 \gg \text{Re}(S) \gg \text{Re}(T) \gg \frac{1}{4\pi^2}$, while the heterotic string loop and sigma model expansions work well in the heterotic string limit: $\text{Re}(S) \gg [4\pi^2 \text{Re}(T)]^3$, $\text{Re}(T) \gg \frac{1}{4\pi^2}$. By varying $\text{Re}(S)$ while keeping $\text{Re}(T)$ fixed, one can smoothly move from the M-theory limit $\epsilon_s \gg 1$ to the heterotic string limit $\epsilon_s \ll 1$ (or vice versa) while keeping $\epsilon_1 \approx \text{Re}(T)/\text{Re}(S)$ and $\epsilon_2 \approx 1/[4\pi^2 \text{Re}(T)]$ small enough. Obviously then the M-theory Kähler potential expanded in ϵ_1 and ϵ_2 remains to be valid over this procedure, and thus is a valid expression of the Kähler potential even in the heterotic string limit. This means that, like the case of the gauge kinetic functions, one can determine the expansion coefficients in (2.8) by matching the heterotic string Kähler potential which can be computed in the string loop and sigma model perturbation theory. Since $\epsilon_1^n \epsilon_2^m \sim$

$\epsilon_s^n \epsilon_\sigma^{m+2n}$, (n, m) -th order in the M-theory expansion corresponds to $(n, m + 2n)$ -th order in the string loop and sigma-model perturbation theory. Thus all the terms in the M-theory expansion have their counterparts in the heterotic string expansion. It appears that the converse is not true in general, for instance the term $\epsilon_s^p \epsilon_\sigma^q$ with $q < 2p$ in the heterotic string expansion does not have its counterpart in the M-theory expansion. However all string one-loop corrections which have been computed so far lead to corrections which scale (relative to the leading terms) as $\epsilon_s \epsilon_\sigma^2$ or $\epsilon_s \epsilon_\sigma^3$, and thus have M-theory counterparts. This leads us to suspect that all the terms that actually appear in the heterotic string expansion have $q \geq 2p$ and thus have their counterparts in the M-theory expansion. Then there will be a complete matching, up to (nonperturbative) corrections which can not be taken into account by the M-theory expansion, of the Kähler potential between the M-theory limit and the heterotic string limit, like the case of the gauge kinetic function and superpotential. Collecting available informations on the coefficients in (2.8), either from the heterotic string analysis or from the direct M-theory analysis (see [11] and references therein) one obtains the following higher order corrections to the leading order Kähler potential in (2.5):

$$\begin{aligned} \delta \hat{K} &= \frac{A_{03}}{[4\pi^2 \text{Re}(T)]^3} \left[1 + \mathcal{O}\left(\frac{1}{4\pi^2 \text{Re}(T)}\right) \right] \\ &+ \frac{A_{11}}{4\pi^2 \text{Re}(S)} \left[1 + \mathcal{O}\left(\frac{1}{4\pi^2 \text{Re}(S)}, \frac{1}{4\pi^2 \text{Re}(T)}\right) \right] \\ \delta Z &= \frac{3}{(T + \bar{T})} \frac{B_{03}}{[4\pi^2 \text{Re}(T)]^3} \left[1 + \mathcal{O}\left(\frac{1}{4\pi^2 \text{Re}(T)}\right) \right] \\ &+ \frac{\beta_O}{2\text{Re}(S)} \left[1 + \mathcal{O}\left(\frac{1}{4\pi^2 \text{Re}(S)}, \frac{1}{4\pi^2 \text{Re}(T)}\right) \right] \end{aligned} \quad (2.9)$$

where A_{03} , A_{11} and B_{03} are of order one.

As a phenomenological application of the M-theory expansion discussed so far, we are going to analyze in subsection 3.2 the soft supersymmetry-breaking terms under the assumption that supersymmetry is spontaneously broken by the auxiliary components F^S and F^T of the moduli superfields S and T . We will see in subsection 3.1 how moduli values of order one are necessary in order to obtain $M_{GUT} \approx 3 \times 10^{16}$ GeV. Clearly, if

$\text{Re}(T)$ is of order one, we are in the M-theory domain with $\epsilon_s \gg 1$. (See (2.3)). One may worry that the M-theory expansion (2.8) would not work in this case since $\epsilon_1 = \text{Re}(T)/\text{Re}(S)$ is of order one also. However as we have noticed, any correction which is n -th order in ϵ_1 accompanies at least $(n - 1)$ -powers of ϵ_2 and thus is suppressed by $(\epsilon_1 \epsilon_2)^{n-1} \approx (\alpha_{GUT}/\pi)^{n-1}$ compared to the order ϵ_1 correction. This allows the M-theory expansion (2.8) to be valid even when ϵ_1 becomes of order one. Obviously if $\text{Re}(T)$ is of order one, only the order ϵ_1 correction to Z , i.e. $\delta Z = \beta_O/2\text{Re}(S)$, can be sizeable. The other corrections are suppressed by either $\epsilon_1 \epsilon_2 \approx 1/4\pi^2 \text{Re}(S)$ or $\epsilon_2^3 \approx 1/[4\pi^2 \text{Re}(T)]^3$ and thus smaller than the leading order results at least by $\mathcal{O}(\frac{\alpha_{GUT}}{\pi})$. Thus we will include only $\delta Z = \beta_O/2\text{Re}(S)$ ($\delta Z = \beta_H/2\text{Re}(S)$ for hidden matter) in the later analysis of soft terms, while ignoring the other corrections to the Kähler potential

Summarizing the above discussion, our starting point of the phenomenological analyses in next sections is the effective supergravity model given by

$$\begin{aligned} K &= -\ln(S + \bar{S}) - 3\ln(T + \bar{T}) \\ &+ \frac{3}{T + \bar{T}} \left(1 + \frac{1}{3}\epsilon_O \right) C_O^p \bar{C}_O^p \\ &+ \frac{3}{T + \bar{T}} \left(1 + \frac{1}{3}\epsilon_H \right) C_H^p \bar{C}_H^p, \end{aligned} \quad (2.10)$$

$$f_O = S + \beta_O T, \quad f_H = S + \beta_H T, \quad (2.11)$$

$$W_O = d_{pqr} C_O^p C_O^q C_O^r, \quad (2.12)$$

with

$$\epsilon_O = \beta_O \frac{T + \bar{T}}{S + \bar{S}}, \quad \epsilon_H = \beta_H \frac{T + \bar{T}}{S + \bar{S}}. \quad (2.13)$$

Notice that the parameter ϵ_O defined above is $\epsilon_O \approx \beta_O \epsilon_1$. Here the superpotential and gauge kinetic functions are exact up to nonperturbative corrections, while there can be small additional perturbative corrections to the Kähler potential which are of order $1/4\pi^2 \text{Re}(S)$ or $1/[4\pi^2 \text{Re}(T)]^3$.

2.2 Universality of soft terms

To carry out an exhaustive analysis of the phenomenology associated to heterotic M-theory compactified on a Calabi-Yau manifold one also should

consider in principle models with more than one single T -modulus. However, models with several moduli T_i has the potential problem of non-universal soft scalar masses [39]. The soft scalar masses are given in general by $m_{p\bar{q}}^2 = m_{3/2}^2 Z_{p\bar{q}} - F^m (\partial_m \partial_{\bar{n}} Z_{p\bar{q}} - Z^{r\bar{s}} \partial_m Z_{p\bar{s}} \partial_{\bar{n}} Z_{r\bar{q}}) \bar{F}^{\bar{n}}$ [40], where $F^m = F^S, F^{T_i}$ in our case, and $Z_{p\bar{q}}$ and $Z^{p\bar{q}}$ denote the Kähler metric and its inverse of the matter fields C^p . For example, in the case of the standard embedding the Kähler metric of the matter fields C^i is given by $Z_{i\bar{j}} = (\partial^2 \hat{K}^T / \partial T_i \partial \bar{T}_j) e^{-\hat{K}^T/3} + \delta Z_{i\bar{j}}(S + \bar{S}, T_k + \bar{T}_k)$, where $\hat{K}^T = -\ln k_{ijk}(T_i + \bar{T}_i)(T_j + \bar{T}_j)(T_k + \bar{T}_k)$ and $\delta Z_{i\bar{j}}$ corresponds to the S -dependent correction in the M-theory expansion (or the string-loop correction). After normalizing the fields to get canonical kinetic terms, although the first piece in $m_{i\bar{j}}^2$ above will lead to universal diagonal soft masses, the second piece will generically induce non-universal contributions, as in the case of the weakly-coupled limit of the heterotic string compactified on a Calabi-Yau [41], due to the presence of the off-diagonal Kähler metric $Z_{i\bar{j}}$ written above. This clearly implies that the scalar mass eigenvalues will be in general non-degenerate. If one ignores $\delta Z_{i\bar{j}}$, the matter Kähler metric is S -independent and, as a consequence, in the dilaton-dominated [42, 43] scenario with $F^{T_i} = 0$ the normalized soft scalar masses are universal as $m_i = m_{3/2}$. However including the S -dependent $\delta Z_{i\bar{j}}$, one generically loses the scalar mass universality even in the dilaton-dominated case [11]. In fact, this was noted in [44] for the string-loop induced³ $\delta Z_{i\bar{j}}$ which is small in the weakly coupled heterotic string limit. The main point here is that in the M-theory limit $\delta Z_{i\bar{j}}$ can be as large as the leading order Kähler metric, and then there can be a large violation of the scalar mass universality even in the dilaton-dominated scenario [11]. An explicit computation of $\delta Z_{i\bar{j}}$ can be found in [46]. Clearly, multimoduli Calabi-Yau models have the potential problem of non-universal soft scalar masses. Of course this can be ameliorated taking into account the low-energy running of the scalar masses [43]. In particular, in the squark case, for gluino masses heavier than

(or of the same order as) the squark masses at the boundary scale, there are large flavour-independent gluino loop contributions which are the dominant source of squark masses. However, to avoid the problem of non-universality from the beginning would be welcome. As pointed out in [13] there is the solution of working with Calabi-Yau spaces with one Kähler modulus T ($h_{1,1} = 1$). Clearly, supersymmetry breaking in the S and/or T direction in this case will give rise to universal soft terms⁴.

Notice that this improvement with respect to the problem of non-universality is not possible in other compactifications. For example, although in most orbifolds the structure of soft scalar masses is simpler due to the existence of diagonal metrics $Z_{p\bar{q}} = \delta_{pq} \prod_i (T_i + \bar{T}_i)^{n_p^i}$, still they show a lack of universality $m_p^2 = m_{3/2}^2 + \sum_i \frac{n_p^i}{(T_i + \bar{T}_i)^2} |F^{T_i}|^2$ [47], due to the modular weight dependence n_p^i [48]. Although the above formulae are valid for the weakly-coupled case, the result about non-universality of soft terms is not modified in the strongly-coupled case [11].

Summarizing the above discussions, due to the constraints that the universality of soft terms impose on effective supergravity models, our starting point for the phenomenological analyses in the next section is the model given by (2.10), (2.11) and (2.12), i.e. we will assume that the standard model arises from heterotic M-theory compactified on a Calabi-Yau manifold with only one modulus field T .

3. Phenomenology of standard and non-standard embedding vacua

Here we will summarize first results found in the literature about the standard and non-standard embedding cases, and then we will discuss in detail the issue of the scales in the theory as well as the pattern of soft terms.

³It is worth noticing that supergravity-loop corrections may also induce non-universality [45].

⁴Although (2,1) complex structure moduli, U_p , may contribute to the matter Kähler metric with some U -dependent metric H_{pq} multiplying the third and forth term in (2.10), they will not spoil the universality of soft terms as long as they do not contribute to supersymmetry breaking, $F^{U_p} = 0$. I thank A. Lukas and D. Waldram for useful discussions about this point.

Let us recall first that the form of the effective action is determined by (2.10), (2.11) and (2.12). This is also true for the non-standard embedding case although there is no requirement that the spin connection be embedded in the gauge connection [24, 25, 26]. Taking into account that the real parts of the gauge kinetic functions in (2.11) multiplied by 4π are the inverse gauge coupling constants α_O and α_H , using (2.1) one can write [3, 5, 24]

$$\alpha_{O,H} = \frac{(4\pi)^{2/3}}{2M_{11}^6 V_{O,H}} , \quad (3.1)$$

with $V_{O,H} = V(1 + \epsilon_{O,H})$ the observable(hidden) sector volume.

On the other hand, using V_O as defined above, the M-theory expression of the four-dimensional Planck scale $M_{Planck}^2 = 16\pi^2 \rho M_{11}^9 < V >$ where $< V >$ is the average volume of the Calabi-Yau space $< V > = (V_O + V_H)/2$, and (2.1) one finds

$$V_O^{-1/6} = \left(\frac{V}{< V >} \right)^{1/2} 3.6 \times 10^{16} \left(\frac{4}{S + \bar{S}} \right)^{1/2} \times \left(\frac{2}{T + \bar{T}} \right)^{1/2} \left(\frac{1}{1 + \epsilon_O} \right)^{1/6} \text{ GeV} , \quad (3.2)$$

which is a very useful formula as we will see below in order to discuss whether or not the GUT scale or smaller scales are obtained in a natural way. In this respect, let us now obtain the connection between the different scales of the theory: the eleven-dimensional Planck mass, M_{11} , the Calabi-Yau compactification scale, $V_O^{-1/6}$, and the orbifold scale, $(\pi\rho)^{-1}$. It is straightforward to obtain from (3.1) the following relation:

$$\frac{M_{11}}{V_O^{-1/6}} = 2 . \quad (3.3)$$

Likewise, using the above expression for M_{Planck} and (3.1) we arrive at

$$\frac{V_O^{-1/6}}{(\pi\rho)^{-1}} = \left(\frac{V}{< V >} \right) \left(\frac{2.7 \times 10^{16} \text{ GeV}}{V_O^{-1/6}} \right)^2 \times 7 (1 + \epsilon_O) . \quad (3.4)$$

Notice that in (3.3) and (3.4) we have already assumed that the gauge group of the observable sector G_O is the one of the standard model or some unification gauge group as $SU(5)$, $SO(10)$

or E_6 , i.e. we are using $(2\pi\alpha_O)^{-1} = 4$ in order to reproduce the LEP data about α_{GUT} (α_O in our notation).

Let us recall at this point that standard and non-standard embedding vacua fulfil the condition (2.7). Thus $\epsilon_O = -\epsilon_H$ in (2.13) implying that the average volume of the Calabi-Yau space turns out to be equal to the lowest order value $< V > = V$ and as a consequence (3.2) and (3.4) simplify. This will not be the case in the presence of five-branes as we will see in the next section.

Due also to eq.(2.7) the following bounds

$$-1 < \epsilon_O < 1 , \quad (3.5)$$

must be fulfilled in order to have positive values for V_O and V_H . Besides, $\epsilon_O > 0$ will imply that V_O be larger than V_H and therefore the gauge coupling of the observable sector will be weaker than the gauge coupling of the hidden sector (see(3.1)). The opposite situation $\epsilon_O < 0$ may be obtained in non-standard embedding models. V_O is now smaller than V_H and therefore the gauge coupling of the observable sector will be stronger than the one of the hidden sector⁵.

Notice that using (2.11) one can write ϵ_O as

$$\epsilon_O = \frac{4 - (S + \bar{S})}{(S + \bar{S})} , \quad (3.6)$$

where $(2\pi\alpha_O)^{-1} = 4$ has been used. Thus with (3.5) and (2.13) one obtains that the dilaton and moduli fields are bounded. In particular,

$$0 < \beta_O(T + \bar{T}) < 2 , \quad 2 < (S + \bar{S}) < 4 , \quad (3.7)$$

for $0 < \epsilon_O < 1$ and

$$\beta_O(T + \bar{T}) < 0 , \quad (S + \bar{S}) > 4 , \quad (3.8)$$

for $-1 < \epsilon_O < 0$. Note that ϵ_O can approach the limit -1 only for very large values of $(S + \bar{S})$ and therefore of $(T + \bar{T})$.

With all these results we can start now the study of scales and soft terms in the theory.

⁵In the context of supersymmetry breaking by gaugino condensation this scenario may have several advantageous features with respect to scenarios with $\epsilon_O > 0$. For a discussion about this point see [25].




Figure 1: $\log M_{11}$, $\log V_O^{-1/6}$ and $\log(\pi\rho)^{-1}$ versus ϵ_O . The straight line corresponds to $M_{GUT} = 3 \times 10^{16}$ GeV.

3.1 Scales

We will discuss first how to obtain $M_{GUT} = 3 \times 10^{16}$ GeV in the four-dimensional effective theory from heterotic M-theory [3, 4] taking into account the higher order corrections studied above to the zeroth-order formulae [27]. On the other hand, we will analyze whether the special limits pointed out in [30], in order to lower the scales of the theory, even with the possibility of obtaining an extra dimension as large as a millimetre, may be obtained in a natural way [27].

Let us concentrate first in the case $\beta_O > 0$, i.e. in the region $0 < \epsilon_O < 1$ in (3.5). Identifying M_{GUT} with $V_O^{-1/6}$ one obtains from (3.4), and (3.3) (recall that $\langle V \rangle = V$): $M_{11} \approx 6 \times 10^{16}$ GeV and $(\pi\rho)^{-1} \approx (2.5 - 5.3) \times 10^{15}$ GeV, i.e. the following pattern $(\pi\rho)^{-1} < V_O^{-1/6} < M_{11}$. On the other hand, to obtain $V_O^{-1/6} = M_{GUT}$ when $\beta_O > 0$ is quite natural. This can be seen from (3.2) since (3.7) implies that $T + \bar{T}$ and $S + \bar{S}$ are essentially of order one. Let us discuss this point in more detail. Using (2.13) and (3.6) it is interesting to write (3.2) as

$$V_O^{-1/6} = 3.6 \times 10^{16} \left(\frac{\beta_O}{2\epsilon_O} \right)^{1/2} \times (1 + \epsilon_O)^{5/6} \text{ GeV} . \quad (3.9)$$

This is shown in Fig. 1 where $V_O^{-1/6}$ versus ϵ_O is plotted. The r.h.s. of the figure ($0 < \epsilon_O < 1$) corresponds to the case $\beta_O > 0$ whereas the l.h.s.

($-1 < \epsilon_O < 0$) corresponds to the case $\beta_O < 0$ that will be analyzed below. For the moment we concentrate on the case $\beta_O > 0$ and, in particular, in Fig. 1 we are showing an example with $\beta_O = 1$. $(\pi\rho)^{-1}$ and M_{11} are also plotted in the figure using (3.4) and (3.3) respectively. Most values of ϵ_O imply $V_O^{-1/6} \approx 5 \times 10^{16}$ which is quite close to the phenomenologically favored value. For example, for $\epsilon_O = 1/4$, which corresponds to $S + \bar{S} = 16/5$ and $T + \bar{T} = 4/5$, we obtain $V_O^{-1/6} = 6.1 \times 10^{16}$ GeV and for the limit $\epsilon_O = 1$ (as discussed in subsection 2.1, the M-theory expansion will work even in this limit), which corresponds to $S + \bar{S} = T + \bar{T} = 2$, we obtain the lowest possible value $V_O^{-1/6} = 4.5 \times 10^{16}$.

These qualitative results can only be modified in the limit $\epsilon_O \rightarrow 0$, i.e. $(T + \bar{T}) \rightarrow 0$, since then $V_O^{-1/6} \rightarrow \infty$. Notice that in this case $(\pi\rho)^{-1} > V_O^{-1/6}$ (see Fig. 1). This limit is not interesting not only because $V_O^{-1/6}$ is too large but also because we are effectively in the weakly-coupled region with a very small orbifold radius.

The results for $\beta_O > 1$ can easily be deduced from the figure and eq.(3.9). For those models we are in the limit of validity if we want to obtain $V_O^{-1/6} = M_{GUT}$. For example, For $\epsilon_O = 1$ with $\beta_O = 4$, $V_O^{-1/6} = 9 \times 10^{16}$ GeV.

Let us finally remark that, from the above discussion, it is straightforward to deduce that large internal dimensions, associated with the radius of the Calabi-Yau and/or the radius of the orbifold, are not allowed.

Let us now study the value of the scales in models with $\beta_O < 0$. We can use again (3.9), but now with $-1 < \epsilon_O < 0$. This is shown in the l.h.s. of the Fig. 1. Unlike the previous models where always $V_O^{-1/6}$ was bigger than M_{GUT} for any $\beta_O > 0$, in these non-standard embedding models M_{GUT} can be obtained. For example in the case shown in the figure, $\beta_O = -1$, with $\epsilon_O = -0.35$ which, using (3.6) and (2.13), corresponds to $S + \bar{S} = 6.15$ and $T + \bar{T} = 2.15$, we obtain $V_O^{-1/6} = 3 \times 10^{16}$ GeV. For other values of β_O this is also possible. Notice that the figure for $V_O^{-1/6}$ will be the same adding the constant $\log |\beta_O|^{1/2}$ and therefore there will be lines, corresponding to $V_O^{-1/6}$, intersecting with the straight line corresponding to M_{GUT} . In this sense, if we want

to obtain models with the phenomenologically favored GUT scale, non-standard embedding models with $\beta_O < 0$ are more compelling than models with $\beta_O > 0$.

On the other hand, for $\beta_O > 0$ we obtained above the lower bound $\approx 10^{16}$ GeV for all scales of the theory (see the r.h.s. of Fig. 1), far away from any direct experimental detection. Now we want to study this issue in cases with $\beta_O < 0$. From (3.9), clearly in the limit $\epsilon_O \rightarrow -1$ we are able to obtain $V_O^{-1/6} \rightarrow 0$ and therefore, given (3.4) also $(\pi\rho)^{-1} \rightarrow 0$ (see the l.h.s. of Fig. 1). Thus to lower the scale $V_O^{-1/6}$ down to the experimental bound (due to Kaluza-Klein excitations) of 1 TeV is possible in this limit. However, this is true only for values of ϵ_O extremely close to -1 . For example, for $\epsilon_O = -0.999999$ which, using (3.6) and (2.13), corresponds to $S + \bar{S} = 4 \times 10^6$ and $T + \bar{T} = 4 \times 10^6 - 4$, we obtain the intermediate scale $V_O^{-1/6} = 2.5 \times 10^{11}$ GeV, i.e. $M_{11} = 5 \times 10^{11}$ GeV, with $(\pi\rho)^{-1} = 3 \times 10^6$ GeV. This is an interesting possibility since an intermediate scale $\approx 10^{11}$ GeV was proposed in [30] in order to solve some phenomenological problems and in [49] in order to solve the M_W/M_{Planck} hierarchy problem⁶. In any case, it is obvious that the smaller the scale the larger the amount of fine-tuning becomes. The experimental lower bound for the scale $V_O^{-1/6}$, 1 TeV, can be obtained with $\epsilon_O = 10^{-16} - 1$, i.e. $S + \bar{S} = 4 \times 10^{16}$ and $T + \bar{T} = 4 \times 10^{16} - 4$. Then one gets $V_O^{-1/6} = 1181.5$ GeV with $(\pi\rho)^{-1} = 3.2 \times 10^{-9}$ GeV. Since only gravity is free to propagate in the orbifold, this extremely small value is not a problem from the experimental point of view. In any case, it is clear that low scales are possible but the fine-tuning needed renders the situation highly unnatural. Another problem related with the limit $\epsilon_O \rightarrow -1$ will be found below when studying soft

⁶For example, for a D3-brane in type I strings where $\frac{M_W}{M_{Planck}} \approx \frac{\alpha_O}{2} \left(\frac{M_c}{M_I}\right)^6$, with a modest input hierarchy between string and compactification scales, $M_I \approx 10^{11}$ GeV and $M_c \approx 10^9$ GeV, one obtains the desired hierarchy $M_W/M_{Planck} \approx 10^{-16}$ without invoking any hierarchically suppressed nonperturbative effect like e.g. gaugino condensation. However, it is worth noticing that those values would imply $\text{Re}(S) = 1/\alpha_O \approx 24$ and $\text{Re}(T) = \frac{1}{\alpha_O} \left(\frac{M_I}{M_c}\right)^4 \approx 10^9$, i.e. one has again a hierarchy problem but now for the vev of the fields that one has to determine dynamically.




Figure 2: $\log M_{11}$, $\log V_O^{-1/6}$ and $\log(\pi\rho)^{-1}$ versus ϵ_O . The straight line corresponds to $M_{GUT} = 3 \times 10^{16}$ GeV.

terms, since $|M|/m_{3/2} \rightarrow \infty$. Thus a extremely small gravitino mass is needed to fine tune the gaugino mass M to the 1 TeV scale in order to avoid the gauge hierarchy problem.

There is a value of β_O which is in principle allowed and has not been analyzed yet. This is the case $\beta_O = 0$. As we will see in a moment, to lower the scales a lot in this context is again possible. Since ϵ_O in (2.13) is vanishing and using (3.6), $S + \bar{S} = 4$, eq. (3.2) can be written as

$$V_O^{-1/6} = 3.6 \times 10^{16} \left(\frac{2}{T + \bar{T}} \right)^{1/2} \text{ GeV}. \quad (3.10)$$

This is plotted in Fig. 2 together with $(\pi\rho)^{-1}$ and M_{11} . We see that the value $V_O^{-1/6} = 3 \times 10^{16}$ GeV is obtained for the reasonable value $T + \bar{T} = 2.88$. On the other hand, the larger $T + \bar{T}$ the smaller $V_O^{-1/6}$ becomes. The lower bound for $V_O^{-1/6}$ is obtained with $T + \bar{T} = 4 \times 10^{19}$ GeV. Then one gets $V_O^{-1/6} = 8 \times 10^6$ GeV and $(\pi\rho)^{-1} = 10^{-13}$ GeV. Smaller values of $V_O^{-1/6}$ are not allowed since experimental results on the force of gravity constrain $(\pi\rho)$ to be less than a millimetre. Thus, although very low scales are allowed for the particular value $\beta_O = 0$, clearly we introduce a hierarchy problem between $S + \bar{S}$ and $T + \bar{T}$.

3.2 Soft terms

Applying the standard (tree level) soft term formulae [40] for the above supergravity model given




Figure 3: Soft parameters in units of $m_{3/2}$ versus θ for different values of ϵ_O . Here M , m and A are the gaugino mass, the scalar mass and the trilinear parameter respectively.

by (2.10), (2.11) and (2.12), one can compute the soft terms straightforwardly⁷ [11]

$$\begin{aligned}
 M &= \frac{\sqrt{3}m_{3/2}}{1 + \epsilon_O} \left(\sin \theta + \frac{1}{\sqrt{3}} \epsilon_O \cos \theta \right), \\
 m^2 &= m_{3/2}^2 - \frac{3m_{3/2}^2}{(3 + \epsilon_O)^2} \left[\epsilon_O (6 + \epsilon_O) \sin^2 \theta \right. \\
 &\quad \left. + (3 + 2\epsilon_O) \cos^2 \theta - 2\sqrt{3}\epsilon_O \sin \theta \cos \theta \right], \\
 A &= -\frac{\sqrt{3}m_{3/2}}{3 + \epsilon_O} \left[(3 - 2\epsilon_O) \sin \theta + \sqrt{3}\epsilon_O \cos \theta \right],
 \end{aligned} \tag{3.11}$$

where $m_{3/2}$ is the gravitino mass, and vanishing cosmological constant and phases are assumed, given the current experimental limits. Here M , m and A denote gaugino masses, scalar masses and trilinear parameters respectively. The bilinear B parameter can be found in [34, 50, 27]. We are using here the parameterization introduced in [43] in order to know what fields, either S or T , play the predominant role in the process of super-

⁷Unlike [18] where only linear terms in ϵ_1 are kept, we keep all contributions to soft terms avoiding accidental cancellations at linear order, e.g. in scalar masses. Higher order terms in (2.10) might modify the higher order contributions but, as argued in subsection 2.1 these terms will be suppressed.

symmetry breaking $F^S = \sqrt{3}m_{3/2}(S + \bar{S}) \sin \theta$, $F^T = m_{3/2}(T + \bar{T}) \cos \theta$.

As mentioned in the introduction, the structure of these soft terms is qualitatively different from that of a Calabi-Yau compactification of the (tree-level) weakly-coupled heterotic string found in [43] which can be recovered from (3.11) by taking the limit $(T + \bar{T}) \ll (S + \bar{S})$, i.e. $\epsilon_O \rightarrow 0$:

$$-A = M = \sqrt{3}m = \sqrt{3}m_{3/2} \sin \theta. \tag{3.12}$$

Clearly the M-theory result (3.11) is more involved due to the additional dependence on ϵ_O . Nevertheless we can simplify the analysis by taking into account the bounds (3.5).

We show in Fig. 3 the dependence on θ of the soft terms M , m , and A in units of the gravitino mass for different values of ϵ_O [11, 31, 27]. Several comments are in order. First of all, some ranges of θ are forbidden by having a negative scalar mass-squared. In the weakly-coupled heterotic string case shown in Fig. 4, the forbidden region vanishes since the squared scalar masses are always positive (see (3.12)). About the possible range of soft terms, the smaller the value of ϵ_O , the larger the range becomes. In the limit $\epsilon_O \rightarrow -1$, $0.3 < |A|/m_{3/2} < 4.58$, $0 < m/m_{3/2} <$




Figure 4: The same as Fig. 3 but for the weakly-coupled heterotic string limit.

2.26 and $|M| \rightarrow \infty$.

In order to discuss the supersymmetric spectra further, it is worth noticing that gaugino masses are in general larger than scalar masses. This implies at low-energy ($\approx M_W$) the following qualitative result [11]: $M_{\tilde{g}} \approx m_{\tilde{q}} > m_{\tilde{l}}$, where $\tilde{g}$ denote the gluino, $\tilde{l}$ all the sleptons and $\tilde{q}$ first and second generation squarks. Other analyses taking into account the details of the electroweak radiative breaking can be found in [34, 35]. Only for values of ϵ_O approaching -1 the opposite situation, scalars heavier than gauginos, may occur. This is for two narrow ranges of values of θ as can be seen in Fig. 3 for $\epsilon_O = -3/5$. Let us remark that $M/m_{3/2}$ and $m/m_{3/2}$ are then very small and therefore $m_{3/2}$ must be large in order to fulfil e.g. the low-energy bounds on gluino masses. In this special limits $m \gg M$ is possible and then $M_{\tilde{g}} < m_{\tilde{q}} \approx m_{\tilde{l}}$ [27].

Notice that in the (tree-level) weakly-coupled heterotic string, the limit $\sin \theta \rightarrow 0$ is not well defined since all M , A , m vanish in that limit. One then has to include the string one-loop corrections (or the sigma-model corrections) to the Kähler potential and gauge kinetic functions which would modify the boundary conditions (3.12). This is similar to what happens in orbifold compactifications where, at the end of the day, scalars are heavier than gauginos due to string loop corrections [43]. This problem is not present in the heterotic M-theory, as can be deduced from Fig. 3, except in models with $\beta_O = 0$, i.e. $\epsilon_O = 0$ and therefore with boundary conditions (3.12).

3.3 Charge and colour breaking

We discussed in subsection 2.2 how effective supergravity models from heterotic M-theory can

be strongly constrained by imposing the (experimental) requirement of universal soft scalar masses, in order to avoid dangerous flavour changing neutral current phenomena. We can go further and impose the (theoretical) constraint of demanding the no existence of low-energy charge and colour breaking minima deeper than the standard vacuum [36]. In this type of analysis, the form of the soft terms is crucial. In the case of the standard embedding, $0 < \epsilon_O < 1$, with soft terms given by (3.11) the restrictions are very strong and the whole parameter space $(m_{3/2}, \theta, \epsilon_O, B)$ turns out to be excluded on these grounds [37, 38]. This is shown in Fig. 5 [38] for a fixed value of $m_{3/2}$ (or, equivalently, of m) with $\epsilon_O = 1$. Then we are left with two independent parameters B and θ . Whereas the black region is excluded because it is not possible to reproduce the experimental mass of the top, the rest is excluded by charge and colour breaking constraints. The small squares indicate regions excluded by the so-called UFB constraints and the circles indicate regions excluded by the so-called CCB constraints. Other values for m and ϵ_O do not modify these conclusions.

Given these dramatic consequences, a way-out must be searched⁸ [51]. The first possibility is to consider the case of the non-standard embedding since although the formulae for the soft

⁸We could accept that we live in a metastable vacuum, provided its lifetime is longer than the present age of the Universe, thus rescuing points in the parameter space [36]. In this sense the constraints found are basically the most conservative ones (in the sense of safe ones).




Figure 5: Excluded regions of the parameter space.

terms are the same (3.11) the parameter space is different: $-1 < \epsilon_O < 0$. Another possible way-out is to consider the presence of five-branes in the vacuum, then the soft terms are different (see (4.2)) and new parameters, as e.g. the goldstino angles θ_n associated with F -terms of the five-branes, enter in the game. Possibly some regions in the parameter space will be allowed. Although now the situation is clearly more model dependent.

It is worth noticing that the situation in the perturbative heterotic string compactified on a Calabi-Yau is basically worst. There the whole parameter space $(m_{3/2}, \theta, B)$ is forbidden [38] and there is no the freedom of playing around with ϵ_O and/or θ_n from five-branes. Only in the limit $\sin \theta \rightarrow 0$, where one has to include loop corrections to the boundary conditions (3.12), small regions might be allowed. At least this is the case of orbifold compactifications with the same boundary conditions (i.e. models where all observable particles have modular weight -1) [38].

4. Vacua with five-branes

In the previous section, we studied the phenomenology of heterotic M-theory vacua obtained through standard and non-standard embedding. Here we want to analyze (non-perturbative) heterotic M-theory vacua due to the presence of five-branes.

4.1 Four-dimensional effective supergravity

As mentioned in the introduction, five-branes are non-perturbative objects, located at points, $x^{11} = x_n (n = 1, \dots, N)$, throughout the orbifold interval. The modifications to the four-dimensional effective action determined by (2.10), (2.11) and (2.12), due to their presence, have recently been investigated by Lukas, Ovrut and Waldram [26, 13]. Basically, they are due to the existence of moduli, Z_n , whose $\text{Re}(Z_n) \equiv z_n = x_n/\pi\rho \in (0, 1)$ are the five-brane positions in the normalized orbifold coordinates. Then, the effective supergravity obtained from heterotic M-theory compactified on a Calabi-Yau manifold in the presence of five-branes is now determined by

$$K = -\ln(S + \bar{S}) - 3\ln(T + \bar{T}) + K_5$$

$$\begin{aligned} & + \frac{3}{T + \bar{T}} \left(1 + \frac{1}{3}e_O\right) H_{pq} C_O^p \bar{C}_O^q, \\ f_O &= S + B_O T, \quad f_H = S + B_H T, \\ W_O &= d_{pqr} C_O^p C_O^q C_O^r, \end{aligned} \quad (4.1)$$

with $e_O = b_O \frac{T + \bar{T}}{S + \bar{S}}$. Here K_5 is the Kähler potential for the five-brane moduli Z_n , H_{pq} is some T -independent metric (see footnote 4) and $b_O = \beta_O + \sum_{n=1}^N (1 - z_n)^2 \beta_n$, $B_O = \beta_O + \sum_{n=1}^N (1 - Z_n)^2 \beta_n$, $B_H = \beta_H + \sum_{n=1}^N (Z_n)^2 \beta_n$, with β_O , β_H the instanton numbers and β_n the five-brane charges. The former, instead of condition (2.7), must fulfil now: $\beta_O + \sum_{n=1}^N \beta_n + \beta_H = 0$.

4.2 Phenomenology

Assuming for simplicity that $\langle Z_n \rangle = \langle z_n \rangle$, i.e. $\langle B_O \rangle = \langle b_O \rangle$, (3.1) is still valid with the modification $\epsilon_{O,H} \rightarrow e_{O,H}$, where $e_H = b_H \frac{T + \bar{T}}{S + \bar{S}}$ with $b_H = \beta_H + \sum_{n=1}^N (z_n)^2 \beta_n$. Following the analysis of section 3 one can write e_O as a function of $S + \bar{S}$ as in (3.6) and therefore the bounds for $S + \bar{S}$ in (3.7) and (3.8) are still valid if $-1 < e_O < 1$ is possible. In fact one can obtain different bounds on e_O depending on the sign of both b_O and b_H [27]. For example, if $b_H \geq 0$ and $b_O \leq 0$, then e_H is positive and e_O negative. Since $V_O = V(1 + e_O)$ must be positive we need $-1 < e_O \leq 0$. Another example is the case $b_H \geq 0$ and $b_O > 0$. Now since e_O is positive V_O will always be positive and therefore the only bound is $e_O > 0$. It is worth noticing that the values $0 < (S + \bar{S}) < 2$, corresponding to $e_O > 1$ are then possible. This was not the case in the absence of five-branes since $\epsilon_O > 1$ was not allowed.

4.2.1 Scales

In the presence of five-branes $V = \langle V \rangle$ as in section 3 is no longer true since $V_{O,H} = V(1 + e_{O,H})$ with $e_O + e_H \neq 0$ in general. Therefore $\frac{V}{\langle V \rangle} = \left[1 + \frac{e_O}{2} \left(1 + \frac{b_H}{b_O}\right)\right]^{-1}$ and the relevant formulae to study the relation between the different scales of the theory are (3.2) and (3.4) with the modification $\epsilon_O \rightarrow e_O$. Notice that (3.3) is not modified. Similarly to the case without five-branes, to obtain $V_O^{-1/6} \approx 3 \times 10^{16}$ GeV when $T + \bar{T}$ and $S + \bar{S}$ are of order one is quite natural. To carry out the numerical analysis we can use




Figure 6: $\log M_{11}$, $\log V_O^{-1/6}$ and $\log(\pi\rho)^{-1}$ versus $\log e_O$. The straight line corresponds to $M_{GUT} = 3 \times 10^{16}$ GeV.

(3.9) with the factor $\frac{V}{\langle V \rangle}$ written above multiplying the r.h.s. and with the modifications $\epsilon_O \rightarrow e_O$, $\beta_O \rightarrow b_O$. Several examples were considered in [27]. Although the qualitative results are similar to those of Fig. 1 with e_O instead of ϵ_O , now the line corresponding to $V_O^{-1/6}$ in the r.h.s. of the figure may intersect the straight line corresponding to the GUT scale. Of course this effect, which is due essentially to the extra factor discussed above, is welcome.

Only in some special limits one may lower the scales. As in the case without five-branes, fine-tuning $e_O \rightarrow -1$ we are able to obtain $V_O^{-1/6}$ as low as we wish. The numerical results will be basically similar to the ones of subsection 3.1. Moreover, as discussed above, $e_O > 1$ is possible in the presence of five-branes and therefore with e_O sufficiently large we may get $V_O^{-1/6}$ very small. This is shown in Fig. 6 for the example $b_O = b_H = 1/2$. For instance, with $\log e_O = 56.1$ the experimental lower bound $(\pi\rho)^{-1} = 10^{-13}$ GeV is obtained for $V_O^{-1/6} = 8 \times 10^6$ GeV, corresponding to $S + \bar{S} = 3.1 \times 10^{-56}$ and $T + \bar{T} = 8$. Clearly we introduce a hierarchy problem.

Finally, the analysis of the special case $b_O = 0$ will be similar to the one of the case $\beta_O = 0$ without five-branes in subsection 3.1. We can use (3.10) with the average volume $[1 + b_H(T + \bar{T})/8]^{-1/2}$ multiplying the r.h.s. Depending on the value of b_H we obtain different results [27]. For example if $b_H > 0$ the results are qualitatively similar to those of Fig. 2, the larger $T + \bar{T}$ the smaller $V_O^{-1/6}$ becomes. However, notice that

now for large T we have a factor $(T + \bar{T})^{-1}$ and then not so large values of $T + \bar{T}$ as in Fig. 2 are needed in order to lower the scales. For example, if $b_H = 1$ then $V_O^{-1/6} = 1$ TeV can be obtained for $T + \bar{T} = 10^{14}$ with the size of the orbifold $(\pi\rho)^{-1} = 5 \times 10^{-12}$ GeV close to its experimental bound of 1 millimetre. In any case, still a large hierarchy between $S + \bar{S}$ and $T + \bar{T}$ is needed.

4.2.2 Soft terms

Let us now concentrate on the computation of soft terms [13, 32, 27]. Due to the possible contribution of several F -terms associated with five-branes, which can have in principle off-diagonal Kähler metrics, the computation of the soft terms turns out to be extremely involved. In order to get an idea of their value and also to study the deviations with respect to the case without five-branes we can do some simplifications. One possibility is to assume that five-branes are present but only the F -terms associated with the dilaton and the modulus contribute to supersymmetry breaking, i.e. $F^{Z_n} = 0$. Then, assuming as before $\langle Z_n \rangle = \langle z_n \rangle$, eq.(3.11) is still valid with e_O instead of ϵ_O . Under these simplifying assumptions, Fig. 3 is also valid in this case since, as discussed above, the range of allowed values of e_O includes those of ϵ_O , i.e. $-1 < e_O < 1$. The relevant difference with respect to the case without five-branes is that now values with $e_O \geq 1$ are allowed. This possibility was studied in [27]. Although the soft terms are qualitatively different from those without five-branes analyzed in Fig. 3, the fact that always scalar masses are smaller than gaugino masses is still true for $e_O \geq 1$. As discussed below (3.12), we will obtain at low-energies, $M_{\tilde{g}} \approx m_{\tilde{q}} > m_{\tilde{l}}$.

Another possibility to simplify the computation of the soft terms is to assume that there is only one five-brane in the model. For example, parameterizing $F^S = \sqrt{3}m_{3/2}(S + \bar{S}) \sin \theta \cos \theta_1$, $F^T = m_{3/2}(T + \bar{T}) \cos \theta \cos \theta_1$, $F^{Z_1} = \sqrt{3}m_{3/2}(\partial_1 \partial_{\bar{1}} K_5)^{-1/2} \sin \theta_1$, where θ_1 is the new goldstino angle associated to the F -term of the five-brane, one obtains [32, 27].

$$M = \frac{\sqrt{3}m_{3/2}}{\left(1 + \frac{B_O T + \bar{B}_O \bar{T}}{S + \bar{S}}\right)} (\sin \theta \cos \theta_1$$

$$\begin{aligned}
& + \frac{1}{\sqrt{3}} B_O \frac{e_O}{b_O} \cos \theta \cos \theta_1 \\
& - \frac{2T}{S + \bar{S}} (1 - Z_1) \beta_1 (\partial_1 \partial_{\bar{1}} K_5)^{-1/2} \sin \theta_1 \Big) , \\
m^2 &= m_{3/2}^2 - \frac{3m_{3/2}^2}{(3 + e_O)^2} \{ e_O (6 + e_O) \sin^2 \theta \\
& \times \cos^2 \theta_1 + (3 + 2e_O) \cos^2 \theta \cos^2 \theta_1 \\
& - 2\sqrt{3} e_O \sin \theta \cos \theta \cos^2 \theta_1 \\
& + (\partial_1 \partial_{\bar{1}} K_5)^{-1} \sin^2 \theta_1 \left((3 + e_O) \beta_1 \frac{e_O}{2b_O} \right. \\
& \left. - \left[(1 - z_1) \beta_1 \frac{e_O}{b_O} \right]^2 \right) + 6(1 - z_1) \beta_1 \frac{e_O}{b_O} \\
& \times (\partial_1 \partial_{\bar{1}} K_5)^{-1/2} \sin \theta \sin \theta_1 \cos \theta_1 \\
& - 2\sqrt{3} (1 - z_1) \beta_1 \frac{e_O}{b_O} (\partial_1 \partial_{\bar{1}} K_5)^{-1/2} \\
& \times \cos \theta \sin \theta_1 \cos \theta_1 \} , \\
A &= -\frac{\sqrt{3} m_{3/2} C}{3 + e_O} \{ (3 - 2e_O) \sin \theta \cos \theta_1 \\
& + \sqrt{3} e_O \cos \theta \cos \theta_1 - (\partial_1 \partial_{\bar{1}} K_5)^{-1/2} \sin \theta_1 \\
& \times \left[(3 + e_O) \partial_1 K_5 + 3(1 - z_1) \beta_1 \frac{e_O}{b_O} \right] \} \quad (4.2)
\end{aligned}$$

The formula for the B parameter can be found in [27]. Unfortunately, the numerical analysis of this simplified case is not straightforward. All soft terms depend not only on the new goldstino angle θ_1 in addition to $m_{3/2}$, θ and e_O , but also on other free parameters. For example, although gaugino masses can be further simplified with the assumption $\langle Z_n \rangle = \langle z_n \rangle$, i.e. $\langle B_O \rangle = \langle \bar{B}_O \rangle = \langle b_O \rangle$ (and therefore $(B_O T + \bar{B}_O \bar{T}) / (S + \bar{S}) = e_O$), and $\langle T \rangle = \langle \bar{T} \rangle$ (and therefore $2T / (S + \bar{S}) = e_O / b_O$), still they have an explicit dependence on z_1 and $\partial_1 \partial_{\bar{1}} K_5$. Notice that, for a given model, β_O and β_1 are known and therefore b_O can be computed once z_1 is fixed. Something similar occurs for the A parameter, where z_1 , $\partial_1 K_5$ and $\partial_1 \partial_{\bar{1}} K_5$ appear explicitly, and for the scalar masses, where z_1 and $\partial_1 \partial_{\bar{1}} K_5$ also appear. Thus in order to compute soft terms when a five-brane is present and contributing to supersymmetry breaking we have to input these values. Fortunately, z_1 is in the range $0 < z_1 < 1$ and, although K_5 is not known, since it depends on z_1 , we expect $\partial_1 K_5$, $\partial_1 \partial_{\bar{1}} K_5 = \mathcal{O}(1)$. So we can consider the following representative case: $z_1 = 1/2$ and $\partial_1 K_5 = \partial_1 \partial_{\bar{1}} K_5 = 1$. Since, still we have to input the value of b_O , we




Figure 7: Soft parameters in units of $m_{3/2}$ versus θ when one five-brane is contributing to supersymmetry breaking.

choose an example with $\beta_O = 1$ and $\beta_1 = -2$ which implies $b_O = b_H = 1/2$. Then all positive values of e_O are allowed. We show in Fig. 7 the soft terms for the value $e_O = 3/5$ with $\theta_1 = \pi/3$. Unlike Fig. 3 without five-branes, we see now a remarkable fact: scalar masses larger than gaugino masses can easily be obtained. This happens not only for narrow ranges of θ . For example, for $e_O = 1/3$ and $\theta_1 = \pi/3$, $\theta \approx 3\pi/2$ one obtains $m/|M| \approx 10$. This result implies a relation of the type $m_{\tilde{l}} \approx m_{\tilde{q}} \approx 3.5 M_{\tilde{g}}$. An exhaustive analysis of other examples can be found in [27].

Acknowledgments

This work has been supported in part by the CICYT, under contract AEN97-1678-E, and the European Union, under TMR contract ERBFMRX-CT96-0090.

References

- [1] For a review, see e.g.: J.H. Schwarz, hep-th/9807135, and references therein.
- [2] P. Hořava and E. Witten, Nucl. Phys. **B460**, 506 (1996); Nucl. Phys. **B475**, 94 (1996).
- [3] E. Witten, Nucl. Phys. **B471**, 135 (1996).
- [4] T. Banks and M. Dine, Nucl. Phys. **B479**, 173 (1996).
- [5] T. Li, J. L. Lopez and D. V. Nanopoulos, Phys. Rev. **D56**, 2602 (1997).
- [6] E. Dudas and C. Grojean, Nucl. Phys. **B507**, 553 (1997).
- [7] H. P. Nilles, M. Olechowski and M. Yamaguchi, Phys. Lett. **B415**, 24 (1997); Nucl. Phys. **B530**, 43 (1998).

- [8] K. Choi, Phys. Rev. **D56**, 6588 (1997).
- [9] H. P. Nilles and S. Stieberger, Nucl. Phys. **B499**, 3 (1997).
- [10] A. Lukas, B. A. Ovrut and D. Waldram, Nucl. Phys. **B532**, 43 (1998).
- [11] K. Choi, H.B. Kim and C. Muñoz, Phys. Rev. **D57**, 7521 (1998).
- [12] H.P. Nilles, Phys. Lett. **B180**, 240 (1986).
- [13] A. Lukas, B.A. Ovrut and D. Waldram, JHEP **9904**, 009 (1999).
- [14] R. Donagi, A. Lukas, B.A. Ovrut and D. Waldram, JHEP **9905**, 015 (1999); hep-th/9901009.
- [15] N. Wyllard, JHEP **9804**, 009 (1998).
- [16] P. Hořava, Phys. Rev. **D54**, 7561 (1996).
- [17] Z. Lalak and S. Thomas, Nucl. Phys. **B515**, 55 (1998).
- [18] A. Lukas, B.A. Ovrut and D. Waldram, Phys. Rev. **D57**, 7529 (1998).
- [19] I. Antoniadis and M. Quiros, Phys. Lett. **B416**, 327 (1998); Nucl. Phys. **B505**, 109 (1997).
- [20] K. Choi, H.B. Kim and H. Kim, Mod. Phys. Lett. **A14**, 125 (1999).
- [21] B. de Carlos, J.A. Casas and C. Muñoz, Phys. Lett. **B299**, 234 (1993).
- [22] B. de Carlos, J. A. Casas and C. Muñoz, Nucl. Phys. **B399**, 623 (1993).
- [23] S. Stieberger, Nucl. Phys. **B541**, 109 (1999).
- [24] K. Benakli, Phys. Lett. **B447**, 51 (1999).
- [25] Z. Lalak, S. Pokorski and S. Thomas, hep-ph/9807503.
- [26] A. Lukas, B.A. Ovrut and D. Waldram, Phys. Rev. **D59**, 106005 (1999).
- [27] D.G. Cerdeño and C. Muñoz, hep-ph/9904444.
- [28] J. Lykken, Phys. Rev. **D54**, 3693 (1996).
- [29] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B436**, 257 (1998).
- [30] K. Benakli, hep-ph/9809582.
- [31] T. Li, hep-ph/9903371.
- [32] T. Kobayashi, J. Kubo and H. Shimabukuro, hep-ph/9904201.
- [33] T. Li, J.L. Lopez and D.V. Nanopoulos, Mod. Phys. Lett. **A12**, 2647 (1997).
- [34] D. Bailin, G.V. Kraniotis and A. Love, Phys. Lett. **B432**, 90 (1998); hep-ph/9812283.
- [35] Y. Kawamura, H.P. Nilles, M. Olechowski and M. Yamaguchi, JHEP **9806**, 008 (1998).
- [36] For a review, see: C. Muñoz, hep-ph/9709329, and references therein.
- [37] S.A. Abel and C.A. Savoy, Phys. Lett. **B444**, 119 (1998).
- [38] J.A. Casas, A. Ibarra and C. Muñoz, hep-ph/9810266, to appear in Nucl. Phys. B.
- [39] For a review, see: C. Muñoz, hep-ph/9710388, and references therein.
- [40] For a review, see: A. Brignole, L.E. Ibanez and C. Muñoz, hep-ph/9707209, and references therein.
- [41] H.B. Kim and C. Muñoz, Z. Phys. **C75**, 367 (1997).
- [42] V.S. Kaplunovsky and J. Louis, Phys. Lett. **B306**, 269 (1993).
- [43] A. Brignole, L. E. Ibanez and C. Muñoz, Nucl. Phys. **B422**, 125 (1994).
- [44] J. Louis and Y. Nir, Nucl. Phys. **B447**, 18 (1995).
- [45] K. Choi, J.S. Lee and C. Muñoz, Phys. Rev. Lett. **80**, 3686 (1998).
- [46] A. Lukas, B.A. Ovrut, K.S. Stelle and D. Waldram, hep-th/9806051.
- [47] A. Brignole, L. E. Ibanez, C. Muñoz and C. Scheich, Z. Phys. **C74**, 157 (1997).
- [48] L.E. Ibañez and D. Lüst, Nucl. Phys. **B382**, 1992 (305).
- [49] C. Burgess, L.E. Ibañez and F. Quevedo, Phys. Lett. **B447**, 257 (1999).
- [50] C. Kokorelis, hep-th/9810187.
- [51] J.A. Casas, A. Ibarra and C. Muñoz, in preparation.